

## Rate of Convergence

The sequences  $w_n = 1/\log_2(n)$ ,  $x_n = 1/n$ ,  $y_n = 1/n^2$ , and  $z_n = 1/2^n$  all converge to 0. How fast? Well,  $w_8 = 1/3$ ,  $x_8 = 1/8$ ,  $y_8 = 1/64$ , and  $z_8 = 1/256$  is true, but doesn't really convey how slow  $w_n \rightarrow 0$ , or how fast  $z_n \rightarrow 0$ . We use functions like these, and more generally  $1/n^p$  and  $1/c^n$  as yardsticks (or benchmarks) with which to compare the speed of convergence of algorithms.

In our context, we are usually trying to compute better and better approximations  $p_n$  to some value  $p$ , and we want to know how fast the error  $e_n = |p - p_n|$  is converging toward 0. It is nice to be able to say  $p_n \rightarrow p$  (our approximations will eventually be close to the answer), but will it take a second or a week? In the example above  $w_{512} > 10^{-1}$ , but  $z_{30} < 10^{-9}$ .

We will use these standard sequences above as benchmarks. If a sequence converges about as slow as the  $w_n$ , we will say it has a logarithmic rate of convergence, but if it goes fast like  $z_n$ , we will say its rate of convergence is exponential.

Here is the formal definition. Think of  $\beta_n$  as one of the sequences given above. If  $\{\beta_n\}$  is a positive sequence converging to 0, then we say that  $p_n \rightarrow p$  with *rate of convergence*  $\beta_n$  if  $\exists N$  and  $k > 0$  such that  $\forall n > N$ ,

$$|p_n - p| \leq k\beta_n.$$

In this case we write  $p = p_n + O(\beta_n)$ .

Now suppose  $p_7 \approx p$ . What does that mean? What about  $p_{18}$ ? Well, if we can write  $p_n = p + O(1/n^3)$ , then we know that  $p_n \rightarrow p$  at least as fast as  $k/n^3 \rightarrow 0$ , for some constant  $k$ , and it is probably safe to say that  $p_{18}$  is about  $(18/7)^3 \approx 17$  times more accurate than is  $p_7$ .

In the same way that we have just measured the error in a discrete setting, we can measure the error associated with a continuous parameter. Suppose  $\lim_{x \rightarrow 0} f(x) = L$ . How fast? We say that  $f(h) \rightarrow L$  as  $h \rightarrow 0$  with *rate of convergence*  $h^q$  if  $\exists \delta$  and  $k > 0$  such that  $\forall |h| \leq \delta$ ,

$$|f(h) - L| \leq kh^q.$$

We write  $f(h) = L + O(h^q)$ . The idea here is that we are approximating  $L$  by  $f(h)$ , and we would like to compare  $|f(h) - L|$  to  $h^q$ , because we have a feeling for how fast  $h^q \rightarrow 0$ .

For example,  $\lim_{x \rightarrow 0} \sin(x) = 0$ . So for small  $x$ ,  $\sin(x) \approx 0$ . But we can do better:

$$\sin(x) = x - \frac{x^3}{3!} + \frac{x^5}{5!} - \cdots,$$

so we can write  $\sin(x) = 0 + O(x)$ , or to give more information  $\sin(x) = x + O(x^3)$ , or even more:  $\sin(x) = x - x^3/6 + O(x^5)$ , etc.