

Lagrange Interpolation on the Roots of Unity

Let n be a positive integer. Define $x_k = e^{2\pi ik/n}$, $k=0:n-1$, where $i = \sqrt{-1}$. Notice that $x_k^n = 1$, and as such these numbers are called the n^{th} roots of unity. They are evenly spaced around the *unit circle* $|z| = 1$ in \mathbb{C} : (De Moivre)

$$x_k = e^{2\pi ik/n} = (\cos(2\pi/n) + i \sin(2\pi/n))^k = \cos(2\pi k/n) + i \sin(2\pi k/n).$$

Now suppose we wish to interpolate the data (x_k, y_k) , $k = 0:n-1$ with a polynomial $p(x) = \sum_{j=0}^{n-1} a_j x^j$ of degree $n-1$ (the Lagrange interpolator).

The Vandermonde view says that p can be determined by solving the system

$$\begin{aligned} Va &= y, \quad \text{where} \\ a &= [a_0, a_1, \dots, a_{n-1}]^T, \quad y = [y_0, y_1, \dots, y_{n-1}]^T \quad \text{and} \\ e_k^T V &= [1, x_k, x_k^2, \dots, x_k^{n-1}]. \end{aligned}$$

Now let's investigate the (Hermitian) matrix $\bar{V}^T V = V^* V = [s_{kj}]$. If $k \neq j$,

$$s_{kj} = \sum_{r=0}^{n-1} \bar{x}_k^r x_j^r = \sum_{r=0}^{n-1} e^{2\pi i(j-k)r/n} = \sum_{r=0}^{n-1} (e^{2\pi i(j-k)/n})^r = \frac{(e^{2\pi i(j-k)/n})^n - 1}{e^{2\pi i(j-k)/n} - 1} = 0,$$

and for $k = j$,

$$s_{jj} = \sum_{r=0}^{n-1} 1^r = n.$$

so $V^* V = nI$. But then $V^* V a = V^* y$, so the coefficients of p are

$$a = V^{-1} y = \left(\frac{1}{n}\right) V^* y.$$

What's the big deal? This is a *discrete Fourier transform* (DFT):

$$p(x) = \sum_{r=0}^{n-1} a_r x^r$$

is a frequency domain polynomial representation of the vector y , the a_r 's are the DFT coefficients (we could write $\hat{y} = \mathcal{F}(y) = a$), and the interpolation conditions $p(x_k) = y_k$ simply say

$$y = \mathcal{F}^{-1}(\mathcal{F}(y)).$$

This DFT, as described here, is simply matrix multiplication by V^* , and requires $O(n^2)$ flops, but using divide-and-conquer to take advantage of the special structure of the roots of unity leads to

$$a = \text{FFT}(y),$$

requiring only $O(n \log(n))$ flops. Polynomials and trigonometric functions famously meet here, and at the Chebyshev polynomials, in some of the most fundamental and elegant of classical mathematics.