

Name: \_\_\_\_\_

(24) 1. Define  $H \equiv H(u) = I - \frac{2}{u^t u} uu^t$ .

(a) Show that  $H(u) = H(\sigma u)$ , for all nonzero scalars  $\sigma$ .

(b) If  $x \in \mathbb{R}^m$  is nonzero, which vector  $u$  should we use so that  $Hx = \beta e_1$ ?

(c) Given  $u \in \mathbb{R}^m$  and  $B \in \mathbb{R}^{m \times n}$ , how many flops are required to compute  $HB$ ?

(d) Let  $u = (3, 4, 0)^t$  and let  $v = (2, 3, 1)^t$ . Compute  $Hv$ .

(25) 2. Let  $A \in \mathbb{R}^{m \times n}$ ,  $m > n$  be full rank.

(a) Describe the thin  $QR$  factorization of  $A$  (not the process, but the resulting output and the properties of  $Q$  and  $R$ ).

(b) Describe the explicit full  $QR$  factorization of  $A$  (not the process, but the resulting output and the properties of  $Q$  and  $R$ ).

(c) The Householder  $QR$  implicit- $Q$  factorization gives a factored  $Q$ . What does this mean?

- (35) 3. Let  $A \in \mathbb{R}^{m \times n}$ ,  $m > n$  and let  $b \in \mathbb{R}^m$ . Let the columns of  $A$  be linearly independent. Consider the least squares problem

$$\min_x \|Ax - b\|_2 \quad (\text{LS}).$$

- (a) Describe the normal equations approach to solving (LS).
- (b) Describe the Gram-Schmidt QR approach to solving (LS).
- (c) Describe the Householder (implicit-Q) QR approach to solving (LS).
- (d) What is the cost (in flops) of each of these methods?
- (e) Describe the conditioning of (LS).

(16) 4. Let  $A = \begin{bmatrix} -3 & 0 \\ 0 & 1 \\ 2 & 1 \end{bmatrix}$ , and let  $b = \begin{bmatrix} 2 \\ 0 \\ 1 \end{bmatrix}$ .

(a) Form the normal equations for these data (you don't have to solve).

(b) Find  $u_1$  for the Householder  $QR$  factorization of  $A$ .

(c) Find  $q_1$ , the first column of the MGS  $QR$  factorization of  $A$ .