

## Least Squares with Gram-Schmidt

Recall the Modified Gram-Schmidt QR factorization:

$$A = QR, \quad \text{where}$$

in exact arithmetic  $Q \in \mathbb{R}^{m \times n}$  satisfies  $Q^T Q = I$  and  $R \in \mathbb{R}^{n \times n}$  is upper triangular with condition number  $\kappa(R) = \kappa(A)$ . The cost is  $2mn^2 + O(mn)$  flops. If  $A$  is overwritten by  $Q$ , then only  $\frac{1}{2}n^2 + O(n)$  extra words of memory are required. If  $\tilde{Q}$  and  $\tilde{R}$  are the computed versions of  $Q$  and  $R$ , then there exists  $\delta A \in \mathbb{R}^{m \times n}$  with  $A + \delta A = \tilde{Q}\tilde{R}$ , where  $\|\delta A\| = \|A\|O(\mu)$ , and  $\|\tilde{Q}^T \tilde{Q} - I\| = \kappa(A)O(\mu)$ .

Now to solve the least squares problem (LS)  $\min_x \|Ax - b\|_2$  we can use back substitution to solve  $Rx = Q^T b$  (to see this substitute  $A = QR$  into the normal equations:  $A^T Ax = A^T b \Rightarrow R^T Q^T QRx = R^T Q^T b \Rightarrow Q^T QRx = Q^T b$ ). Notice that when we write  $Rx = Q^T b$  we are assuming  $Q^T Q = I$ . This is true in exact arithmetic, but the result above says that in finite precision, the orthogonality of  $Q$  depends on  $\kappa(A)$ . Unfortunately, this – combined with the conditioning of  $R$  – gives a  $[\kappa(A)]^2$  factor in the backward error for  $x$ . Here we will show how to avoid this to get a backward error result for MGS which is equivalent to that of the Householder QR applied to (LS).

Let the MGS QR factorization of  $[A, b]$  be written as

$$\begin{bmatrix} A & b \end{bmatrix} = \begin{bmatrix} Q & q \end{bmatrix} \begin{bmatrix} R & r \\ 0 & \rho \end{bmatrix},$$

where  $q^T Q = 0$  ( $q$  is just  $q_{n+1}$ ). Note that  $A = QR$  and  $b = Qr + \rho q$ . Then  $Q^T b = Q^T(Qr + \rho q) = r$ , and (LS) is solved by backward substitution:  $Rx = r$ . Just for fun, show (both algebraically and geometrically) that  $\rho = \min_x \|Ax - b\|_2$ .

If you already had the MGS factorization  $A = QR$  before  $b$  arrived, no worries. The partitioning above is just a nice way to package the algebra of one more MGS step:

```
w = b
for i=1:n,
    r(i) = Q(:,i)'*w
    w = w - r(i)*Q(:,i)
end
rho = norm(w);  q = w/rho;
```

Now do you really think that this removes the  $\kappa(A)$  factor which came from the orthogonality errors in  $Q$ ? Why should it? I applaud you for your skepticism. The answer lies in the (substantial) differences in behavior between MGS and CGS. Explicitly computing  $Q^T b$  as  $r_c = Q^T * b$  is the CGS way, but MGS (the loop above) adapts to the errors made in each inner product, giving an  $r$  which has, (to the extent that it can), “accounted for” nonorthogonality in the columns of  $Q$ .